

## 業 績

### I 原著

Kawamoto R, Kohara K, Katoh T, Kusunoki T, Ohtsuka N, Abe M, Kumagi T, Miki T. Effect of weight loss on central systolic blood pressure in elderly community-dwelling persons.  
Hypertens Res. 2014; 37: 933-938. [IF:2.936]

Kawamoto R, Kohara K, Katoh T, Kusunoki T, Ohtsuka N, Abe M, Kumagi T, Miki T. Brachial-ankle pulse wave velocity is a predictor of walking distance in community-dwelling adults.  
Aging Clin Exp Res. 2014 Jul 19. [IF:1.136]

Kawamoto R, Kohara K, Katoh T, Kusunoki T, Ohtsuka N, Abe M, Kumagi T, Miki T. Changes in oxidized low-density lipoprotein cholesterol are associated with changes in handgrip strength in Japanese community-dwelling persons.  
Endocrine. 2014 Jul 27. [IF:3.527]

Kawamoto R, Ninomiya D, Hasegawa Y, Kasai Y, Kusunoki T, Ohtsuka N, Kumagi T, Abe M. Mildly elevated serum bilirubin levels are negatively associated with carotid atherosclerosis among elderly persons.  
PLoS One. 2014 Dec 5; 9: e114281. [IF:3.534]

Kawamoto R, Ninomiya D, Hasegawa Y, Kasai Y, Kusunoki T, Ohtsuka N, Kumagi T. Association between serum bilirubin and estimated glomerular filtration rate among elderly persons.  
PLoS One. 2014 Dec 16; 9: e115294. [IF:3.534]

Lammers WJ, van Buuren HR, Hirschfield GM, Janssen HLA, Invernizzi P, Mason AL, Talwalkar J, Ponsioen CY, Floreani A, Corpechot C, Mayo MJ, Battezzati PM, Parés A, Nevens F, Burroughs AK, Kowdley KV, Trivedi PJ, Kumagi T, Cheung A, Lleo A, Imam M, Boonstra K, Cazzagon N, Franceschet I, Poupon R, Caballeria L, Pieri G, Kanwar P, Lindor KD, Hansen BE – on behalf of the Global PBC Study Group.  
Levels of alkaline phosphatase and bilirubin values are surrogate endpoints of outcome for patients with primary biliary cirrhosis – an international follow-up study.  
Gastroenterology. 2014; 147: 1338-1349. [13.926]

Trivedi P, Kumagi T, Al-Harthy N, Coltescu C, Ward S, Cheung AC, Hirschfield GM.  
Good maternal and fetal outcomes in pregnancy associated with primary biliary cirrhosis.  
*Clin Gastroenterol Hepatol.* 2014; 12: 1179-1185. [6.534]

Trivedi P, Bruns T, Cheung AC, Li KK, Kumagi T, Shah H, Corbett C, Al-Harthy N, Acarsu U, Coltescu C, Tripathi D, Neuberger J, Janssen H, Hirschfield GM  
Optimising risk stratification in primary biliary cirrhosis: AST/platelet ratio index predicts outcome independent of ursodeoxycholic acid response.  
*J Hepatol.* 2014; 60: 1249–1258. [10.401]

Hirooka M, Ochi H, Koizumi Y, Tokumoto Y, Hiraoka A, Kumagi T, Abe M, Tanaka H, Hiasa Y  
Local recurrence of hepatocellular carcinoma in the tumor blood drainage area following radiofrequency ablation.  
*Mol Clin Oncol.* 2014; 2: 182-186. [N/A]

Miyake T, Kumagi T, Furukawa S, Hirooka M, Kawasaki K, Koizumi M, Todo Y, Yamamoto S, Abe M, Kitai K, Matsuura B, Hiasa Y.  
Hyperuricemia is a risk factor for the onset of impaired fasting glucose in men with a high plasma glucose level: a community-based study.  
*PLoS One.* 2014; 9: e107882. [3.534]

Furukawa S, Yamamoto S, Maruyama K, Miyake T, Ueda T, Niiya T, Takatoshi S, Torisu M, Kumagi T, Sakai T, Minami H, Miyaoka H, Matsuura B, Hiasa Y, Onji M, Tanigawa T.  
Association between subclinical hypothyroidism and diabetic nephropathy in patients with type 2 diabetes mellitus.  
*Endocrine J.* 2014; 61; 1011-1018. [2.019]

## II 症例報告

Watanabe T, Tokumoto Y, Hirooka M, Koizumi Y, Tada F, Ochi H, Abe M, Kumagi T, Ikeda Y, Matsuura B, Takada K, Hiasa Y.  
An HBV-HIV co-infected patient treated with tenofovir-based therapy who achieved HBs antigen/antibody seroconversion.  
*Intern Med.* 2014; 53: 1343-1346. [0.967]

Ohno Y, Kumagi T, Kuroda T, Koizumi M, Azemoto N, Yamanishi H, Oda M, Hirooka M, Abe

M, Ikeda Y, Matsuura B, Onji M, Soga Y, Mizuno Y, Sugita A, Hiasa Y.  
Signet-ring cell carcinoma of the gallbladder complicated with pulmonary tumor thrombotic microangiopathy.  
Intern Med. 2014; 53:1125-1129. [0.967]

## V 國際学会発表

### **European Association for the Study of the Liver 2014 (2014.4.9-13 London, UK)**

Effective stratification of hepatocellular carcinoma risk in primary biliary cirrhosis: results of a multi-centre international study.

Trivedi PJ, Lammers WJ, van Buuren HR, Janssen HL, Invernizzi P, Battezzati PM, Floreani A, Parés A, Ponsioen CY, Corpechot C, Poupon R, Mayo MJ, Talwalkar J, Burroughs AK, Nevens F, Mason AL, Bruns T, Li K-K, Kowdley KV, Leeman M, Kumagi T, Cheung A, Lleo A, Cazzagon N, Franceschet I, Caballeria L, Boonstra K, de Vries EMG, Imam M, Pieri G, Kanwar P, Lindor KD, Hansen BE, Hirschfield GM; the Global PBC Study Group.

### **65th Annual Meeting of American Association for the Study of the Liver (2014.11. 7 - 11. Boston, USA)**

Pancreatic congestion in liver cirrhosis correlates with impaired insulin secretion

Kuroda T, Hirooka M, Koizumi M, Ochi H, Hisano Y, Bando K, Matsuura B, Kumagi T, Hiasa Y

New model to identify UDCA-treated primary biliary cirrhosis patients in need of additional therapy. Results of an international follow-up study of 4119 patients

Lammers WJ, van Buuren HR, Ponsioen CY, Janssen HL, Floreani A, Hirschfield GM, Corpechot C, Mayo MJ, Invernizzi P, Battezzati PM, Pares A, Nevens F, Burroughs AK, Mason A, Kowdley KV, Imam M, Boonstra K, Cheung AC, Kumagi T, Cazzagon N, Franceschet I, Trivedi PJ, Poupon R, Lleo A, Caballeria L, Pieri G, Lindor KD, Hansen BE

Sub-stratification of hepatocellular carcinoma risk in men with primary biliary cirrhosis: results of an international multicenter study

Trivedi PJ, Lammers WJ, van Buuren HR, Floreani A, Pares A, Cheung AC, Ponsioen CY, Corpechot C, Mayo MJ, Invernizzi P, Battezzati PM, Nevens F, Mason A, Kowdley KV, Li KK, Bruns T, Imam M, Kumagi T, Cazzagon N, Franceschet I,

Caballeria L, Boonstra K, Poupon R, Lleo A, Lindor KD, Janssen HL, Hansen BE,  
Hirschfield GM