

Terakado N, Shintani S, Nakahara Y, Yano J, Hino S, Hamakawa H.

Conservative treatment of prolonged bilateral mandibular dislocation with the help of an intermaxillary fixation screw.

Br J Oral Maxillofac Surg. 44(1):62-63, 2006.2

Shintani S, Li C, Mihara M, Klosek SK, Terakado N, Hino S, Hamakawa H.

Anti-tumor effect of radiation response by combined treatment with angiogenesis inhibitor, TNP-470, in oral squamous cell carcinoma.

Oral Oncol. 42(1):66-72, 2006.1

Elsheikh MN, Rinaldo A, Hamakawa H, Mahfouz ME, Rodrigo JP, Brennan J, Devaney KO, Grandis JR, Ferlito A.

The importance of molecular analysis in detecting cervical lymph node metastasis in head and neck squamous cell carcinoma.

Head Neck. 28(9):842-849, 2006.9

Hara S, Nakashiro K, Klosek SK, Ishikawa T, Shintani S, Hamakawa H.

Hypoxia enhances c-Met/HGF receptor expression and signaling by activating HIF-1 α in human salivary gland cancer cells.

Oral Oncol. 42(6):593-598, 2006.7

Nakahara Y, Shintani S, Mihara M, Hino S, Hamakawa H.

Detection of p16 promoter methylation in the serum of oral cancer patients.

Int J Oral Maxillofac Surg. 35(4):362-365, 2006.4

Ishikawa T, Nakashiro K, Hara S, Klosek SK, Li C, Shintani S, Hamakawa H.

CXCR4 expression is associated with lymph-node metastasis of oral squamous cell carcinoma.

Int J Oncol. 28(1):61-66, 2006.1

Sugimori E, Shintani S, Ishikawa K, Hamakawa H.

Effects of apatite foam combined with platelet-rich plasma on regeneration of bone defects.

Dent Mater J. 25(3): 591-596, 2006.9

Maeda A, Nakashiro K, Hara S, Sasaki T, Miwa Y, Tanji N, Yokoyama M, Hamakawa H, Oyasu R.

Inactivation of AR activates HGF/c-Met system in human prostatic carcinoma cells.
Biochem Biophys Res Commun. 347(4):1158-1165, 2006.9

Isokane M, Sumida T, Okuhira T, Shintani S, Hamakawa H.
Surface osteosarcoma: 2 case reports.
Am J Otolaryngol.27(5):349-352, 2006.9-10

Shintani S, Zhang T, Aslam A, Sebastian K, Yoshimura T, Hamakawa H.
P53-dependent radiosensitizing effects of Hsp90 inhibitor
17-Allylamino-17-demethoxygeldanamycin on human oral squamous cell carcinoma cell
lines.
Int J Oncol. 29(5):1111-1117, 2006.11

Nakaya K, Yamagata HD, Arita N, Nakashiro KI, Nose M, Miki T, Hamakawa H.
Identification of homozygous deletions of tumor suppressor gene FAT in oral cancer
using CGH-array.
Oncogene.26(36):5300-5308, 2007.8

Yoshimura T, Sumida T, Liu S, Onishi A, Shintani S, Desprez PY, Hamakawa H.
Growth inhibition of human salivary gland tumor cells by introduction of progesterone
(Pg) receptor and Pg treatment.
Endocr Relat Cancer. 14(4):1107-1116, 2007.12

Murase R, Abe Y, Takeuchi T, Nabeta M, Imai Y, Kamei Y, Lisa Kagawa-Miki, Ueda N,
Sumida T, Hamakawa H, Kito K
Serum autoantibody to sideroflexin 3 as a novel tumor marker for oral squamous cell
carcinoma.
Proteomics Clin Appl. 2(4): 517-527, 2008.4

Hara S, Nakashiro K, Goda H, Hamakawa H
Role of Akt isoforms in HGF-induced invasive growth of human salivary gland cancer
cells.
Biochem Biophys Res Commun.370(1): 123-128, 2008.5

Hamakawa H, Nakashiro K, Sumida T, Shintani S, JN Myers, RP Takes, A Rinaldo, A
Ferlito

Basic evidences of molecular targeted therapy for oral cancer and salivary gland cancer.
Head Neck. 30(6): 800-809, 2008.6

M Isokane, M Hieda, S Hirakawa, M Shudou, K Nakashiro, K Hashimoto, H Hamakawa,
Higashiyama S

Plasma-membrane-anchored growth factor pro-amphiregulin binds A-type lamin and
regulates global transcription.

J Cell Sci. 121(21): 3608-3618, 2008.11

A Ferlito, A Rinaldo, KO Devaney, Nakashiro K, Hamakawa H

Detection of lymph node micrometastases in patients with squamous carcinoma of the
head and neck.

Eur Arch Otorhinolaryngol. 265 : 1147-1153, 2008.10

Sebastian K, Nakashiro K, Hara S, Goda H, Hamakawa H

Stat3 as a molecular target in RNA interference-based treatment of oral squamous cell
carcinoma.

Oncol Rep. 20(4): 873-878, 2008.10

Klosek SK, Nakashiro K, Hara S, Goda H, Hasegawa H, Hamakawa H

CD151 regulates HGF-stimulated morphogenesis of human breast cancer cells.

Biochem Biophys Res Commun. 379(4): 1097-1100, 2009.2

Ishikawa T, Nakashiro K, Klosek SK, Goda H, Hara S, Uchida D, Hamakawa H

Hypoxia enhances CXCR4 expression by activating HIF-1 in oral squamous cell
carcinoma.

Oncol Rep. 21(3): 707-12, 2009.3

Hamada K, Zhang T, Desaki J, Nakashiro K, Itoh H, Tani K, Koyama Y, Hamakawa H

Carrier cell-mediated cell lysis of squamous cell carcinoma cells by squamous cell
carcinoma antigen 1 promoter-driven oncolytic adenovirus.

J Gene Med. 12(6): 545-54, 2010.6